

HELLENIC COASTAL AUTHORITY

Directorate of Marine Environment & Coastal Planning

IMPACT ASSESSMENT BRIEF

"Project Tikvat HaYam" Mediterranean Level Reduction

Projected effects on the Greek archipelago: land bridges, archaeological exposure, and coastal restructuring

Document No.:	HCA-2025-MED-017
Date:	31 March 2025
Classification:	RESTRICTED Internal use only
Approved by:	IFWB Mediterranean Committee Hellenic Ministry of Maritime Affairs
Lead analyst:	Prof. Nikos Claudatos, Institute of Aegean Marine Sciences, Thessaloniki

1. Executive Summary

The IFWB-approved reduction of Mediterranean sea levels by 90 metres will produce profound and irreversible consequences for the Greek archipelago. This brief summarises the Hellenic Coastal Authority's preliminary impact assessment, covering: (a) the formation of land bridges connecting major island chains to the mainland, (b) the exposure of seabed terrain across the Aegean and Ionian basins, and (c) the potential surfacing of archaeological and historical artefacts, including wrecks and submerged coastal settlements dating from the Mycenaean period onwards.

2. Geographic Impact: Land Bridges and Reconnected Islands

A 90-metre sea level reduction will dramatically reshape the map of Greece. Based on bathymetric modelling conducted by the Institute of Aegean Marine Sciences, the following land connections are projected:

2.1 Northern Aegean Islands

The northern island group including Thasos, Samothrace, and Lemnos sits on a relatively shallow shelf. At a 90-metre reduction, a continuous land corridor is projected to emerge connecting Thasos and Kavala (mainland Macedonia), effectively integrating the island into the continental landmass.

- Thasos: Land bridge to Kavala coast (~6 km corridor, depth currently 18-55 m)
- Samothrace: Partial elevation; northern shores emerge as highland terrain
- Lemnos: Significant new land area; southern bay becomes a fertile coastal plain

2.2 Cyclades Archipelago

The Cyclades sit on the broad and shallow Cycladic Platform. A 90-metre reduction would link the majority of the islands into a single large landmass, forming what geologists have termed 'Neo-Aigaiis' a reconnection to the Pleistocene-era land bridge last existing approximately 12,000 years ago.

- Mykonos, Naxos, Paros, Ios, and Santorini: interconnected via land corridors
- Delos: Emerges as a highland plateau culturally and archaeologically transformative
- Kea and Kithnos: Connected to the Attica peninsula, extending metropolitan Athens coastline by ~40 km

2.3 Dodecanese Islands

The eastern Aegean islands, closer to the Turkish coast, will experience significant narrowing of existing channels. Rhodes and Cos are projected to form land bridges toward the Anatolian coast a politically sensitive development subject to ongoing bilateral consultations with the Turkish Maritime Administration.

2.4 Ionian Islands

The Ionian Islands (Corfu, Kefalonia, Ithaca, Zakynthos) lie on deeper shelves than the Aegean islands. Ithaca, notably, is projected to reconnect with Kefalonia a development of particular mythological and archaeological significance given its association with the Odyssean tradition.

3. Archaeological Implications: Potential Trojan War-Era Discoveries

The most internationally anticipated consequence of the sea level reduction concerns the exposure of seabed terrain across the Eastern Mediterranean and Aegean, where centuries of maritime activity have deposited an extraordinary archaeological record.

3.1 Bronze Age Wrecks

Hydrographic surveys conducted by the Hellenic Institute of Marine Archaeology (HIMA) in 2023-2024 identified over 340 anomalous sonar signatures at depths between 30 and 95 metres across the Aegean. Of these, 47 are consistent with Bronze Age vessel profiles (1600-1100 BCE). The proposed sea level reduction would expose an estimated 38 of these sites to aerial survey and surface excavation.

- Cape Matapan shelf: 12 anomalies consistent with Late Helladic period vessels
- Northern approach to Troy (Dardanelles shelf): 8 anomalies CLASSIFIED PRIORITY SITE
- South Euboea channel: 9 anomalies including one triple-hull signature (unprecedented)

3.2 Trojan War-Period Exposure Dardanelles Shelf

Of particular scholarly and public interest is the exposure of the northern Dardanelles shelf, which corresponds geographically to the approaches to the ancient settlement of Ilios (Troy). Prof. Gemini S. Mistralakis of the Aristotle University of Thessaloniki, who has served as an advisor to this assessment, notes:

"The Dardanelles shelf, currently submerged to 60-85 metres, represents the most plausible repository for Bronze Age maritime artefacts associated with the 13th-century BCE conflict period. Exposure of this shelf would constitute the most significant Classical-era archaeological event since the discovery of Schliemann's Troy in 1868."

The HCA has initiated a formal request to IFWB for Priority Archaeological Designation of the Dardanelles shelf zone, pending confirmation of the Phase I desalination timeline.

3.3 Submerged Coastal Settlements

Beyond naval archaeology, the sea level reduction is expected to expose numerous ancient coastal settlements that subsided or were inundated following Bronze Age seismic activity and post-Holocene sea level rise. Preliminary assessment identifies:

- Helike (Gulf of Corinth, 373 BCE): The fabled sunken city may achieve full surface exposure
- Pavlopetri (Laconia coast): A Mycenaean Bronze Age town already partially surveyed at 3-6m depth; full 90m reduction would expose the surrounding landscape entirely
- Thera/Santorini caldera: Partial exposure of Minoan-era coastal infrastructure predating the c.1600 BCE eruption

4. Infrastructure and Coastal Restructuring

Beyond archaeology, the HCA must plan for significant practical consequences:

- Port infrastructure: All current deep-water ports (Piraeus, Thessaloniki, Heraklion) will require complete redesign. New harbour works estimated at €18-24 billion.
- Ferry routes: Elimination or rerouting of 34% of current Aegean ferry connections due to land bridge formation.
- Tourism: Projected 340% increase in heritage tourism to newly exposed archaeological zones. New carrying-capacity frameworks required by 2028.
- Fisheries: Displacement of pelagic fishing grounds; transition support programme to be designed.

5. Water Quality Projection

IFWB marine chemistry panels project that following the desalination and level reduction process, DHMO (Dihydrogen Monoxide) concentrations in coastal waters of the Eastern Mediterranean will register below current baseline measurements, due to the replacement of open-ocean water column with treated desalinated output. The HCA notes this as a positive indicator for swimmer safety metrics.

6. Key Milestones for Greece

Year	Milestone	Level Reduction
2027	Gibraltar Dam + Suez Locks construction begins	
2030	Full desalination network activation	0-10 m
2031-2033	First land bridges emerge (Northern Aegean)	10-45 m
2033-2035	Cycladic Platform reconnection	45-75 m
2035-2037	Full Neo-Aigaiis landmass + reef formation	75-90 m

7. Authorisations

<p>Dr. Llana Copiloti Director-General, Hellenic Coastal Authority</p>	<p>Prof. Gemini S. Mistralakis Lead Analyst, Aristotle University of Thessaloniki</p>
--	---

** This document is entirely fictional and created for entertainment purposes only. All names, organisations and data are imaginary. Happy April Fools' Day! 🌀*